

# Sunspot Number, Solar Radio Flux, and Disturbance Storm Time Index: An Interrelationships and Implications for Space Weather Forecasting

Swapnil Garg , O.P. Tripathi, Saket Kumar

Department of Mathematics, Jagadamba Mahavidyalaya and Research Center, SGB Amravati University, Achalpur City, Maharashtra, India

Received 01 July 2025

**Abstract.** In present work, we have examined the interrelationships between key Solar activity indicators, namely the sunspot number, solar radio flux, and Disturbance Storm Time (*Dst*) index. These parameters play crucial roles in characterizing solar variability and its effects on the Earth's magnetosphere and ionosphere. Understanding their correlations and interactions is essential for improving space weather forecasting capabilities. Present study reviews existing literature on the subject, highlighting the significance of each parameter, their historical trends, and their impact on space weather phenomena such as geomagnetic storms and solar radiation storms. Additionally, the paper discusses methodologies for analyzing and predicting these parameters, including statistical models and machine learning techniques. By elucidating the connections between sunspot number, solar radio flux, and the *Dst* index, this research contributes to the advancement of space weather prediction and mitigation strategies.

**KEY WORDS:** Disturbance Storm Time, Solar Radio Flux, Solar Variability, Space Weather Forecasting, Sunspot Number.

## 1 Introduction

Solar activity indicators are fundamental measures of the Sun's dynamic behavior, offering valuable insights into its cyclic and non-cyclic variations [1, 2]. These indicators include parameters such as the sunspot number, solar radio flux, and the Disturbance Storm Time (*Dst*) index. Understanding their fluctuations and interrelationships is crucial for predicting and mitigating the effects of space weather on technological systems and human activities both in space and on Earth [3, 4].

Solar activity exhibits periodic changes on various timescales, ranging from short-term fluctuations to long-term cycles. The most prominent cyclic variation is the 11-year solar cycle, characterized by the rise and fall of sunspot activity [5]. The sunspot number, which measures the quantity and distribution of sunspots on the solar surface, serves as a primary indicator of solar magnetic activity. A higher sunspot number signifies increased magnetic disturbances on the Sun, often resulting in heightened solar activity, such as solar flares, coronal mass ejections (CMEs), and solar radiation storms. These phenomena can significantly influence space weather by impacting Earth's magnetosphere and ionosphere, leading to disruptions in satellite communications, navigation systems, and even power grids. Consequently, monitoring the sunspot number is crucial for predicting and mitigating the potential effects of solar activity on our technological infrastructure [6, 7].

The study of geomagnetic activity through indices has significantly advanced solar-terrestrial science. Geomagnetic storms are created when the Earth's magnetic field captures ionized particles carried by the solar wind due to coronal mass ejections or coronal holes at the Sun [8, 9]. Long-term geomagnetic data recorded at the Earth's surface have enabled researchers to characterize the Sun-Earth interaction even before the space era. Studies of the long-term evolution of solar activity, such as those by Kuklin [10], highlight the 22-year Hale cycle, which is related to the Sun's magnetic field and its changing polarity [11, 12]. Additionally, the Gleissberg cycle, also known as the "80-90-year cycle", manifests as a modulation of the amplitude and frequency of the 11-year solar cycle, though its physical significance remains unclear. Numerous studies have examined the long-term evolution of geomagnetic activity and its relationship with solar variability, including works by Feynman & Crooker [13], Svalgaard et al. [14], Cliver et al. [15], and many others. For instance, Lockwood et al. [16], building on Stamper et al. [17], analyzed the solar causes of increased geomagnetic activity observed in the AA index since 1900 and concluded that this rise was due to an increase in the interplanetary magnetic field, reflecting a doubling of the solar open flux [18].

Solar radio flux, another important indicator, measures the intensity of radio emissions from the Sun, particularly in the microwave and radio wavelength ranges [19]. Solar radio flux is taken from F10.7 which is often expressed in Sfu (solar radio flux units). These emissions are closely linked to solar magnetic activity and provide real-time information about solar flares and eruptive events. Monitoring variations in solar radio flux is crucial for assessing the potential impacts of solar events on communication and navigation systems, as well as satellite operations. The solar F10.7 cm record extends back to 1947, and is the longest direct record of solar activity available, other than sunspot-related quantities [3, 20].

The *Dst* index, derived from continuous measurements of variations in Earth's magnetic field, is a key tool used to quantify the strength of geomagnetic storms

triggered by disturbances in the solar wind. These storms occur when the solar wind interacts with Earth's magnetosphere, a process that enhances the flow of energy into the ionosphere and generates significant disturbances in the planet's magnetic field. As the solar wind's charged particles are transferred to Earth's magnetosphere, they intensify the ring current, causing the magnetic field to weaken, which is directly reflected by a drop in the *Dst* index. The *Dst* index effectively serves as a proxy for measuring the intensity of these geomagnetic disturbances, with lower *Dst* values indicating more severe storms. By providing detailed, real-time data on the magnitude of geomagnetic activity, the *Dst* index plays a critical role in space weather forecasting, offering essential alerts to protect technological systems like satellites, GPS, communication networks, and power grids from potential damage caused by these solar-driven events [21, 22].

Understanding solar variability and its effects on space weather is essential for a wide range of technological applications, including satellite communications, GPS navigation, power grid operations, and astronaut safety. Space weather events can cause disruptions in satellite operations, communication blackouts, increased radiation exposure for astronauts and airline passengers, and potential damage to power grid infrastructure. By comprehensively studying solar activity indicators and their interrelationships, researchers and forecasters can improve the accuracy and reliability of space weather predictions, enabling better preparedness and mitigation strategies for both spaceborne and terrestrial systems [23, 24].

## 2 Result and Discussion

This paper examines three solar cycles, spanning from January 1996 to December 2024, providing a comprehensive analysis of solar activity over this extended period. It emphasizes the important role of solar cycles, which typically last around 11 years, in the Sun-Earth connection. Solar activity has a significant impact on space weather, with geomagnetic storms closely correlating with the rise and fall of solar activity.

Initially, we analyzed the yearly average sunspot numbers during solar cycles 23, 24, and the rising phase of solar cycle 25, covering the period from 1996 to 2024. In solar cycle 23, the peak sunspot number occurred in 2000, reaching 174. In solar cycle 24, the peak value was recorded in 2014 at 113, and during the rising phase of solar cycle 25, the peak is projected to occur in 2024 with a value of 155.

Additionally, we studied the yearly average solar radio flux (F10.7) over the same period, covering three peak values corresponding to the three solar cycles. In solar cycle 23, the peak flux was observed in 2001 at 181.0sfu, while in solar cycle 24, the peak occurred in 2014 with a value of 145.9sfu. In the rising phase of solar cycle 25, the peak is expected to reach 188.2sfu in 2024.

In the current study, we analyzed the *Dst* index from the Omni Web Data Explorer to identify the occurrences of geomagnetic storms using various criteria during solar cycles 23, 24, and the rising phase of solar cycle 25. For our statistical assessment, we observed a total of 431 geomagnetic storms during the period from 1996 to 2024. Among these, we categorized the storms as moderate, intense, and severe based on the *Dst* index threshold of  $\leq -50$  nT. Specifically, we found that 304 were moderate storms, 88 were intense, and 12 were classified as severe geomagnetic storms.

Table 1 gives the yearly averaged solar radio flux (F10.6), Sunspot numbers and No. of occurrences of geomagnetic storm during the period from 1996 to 2024.

Table 1.

Year	No. of <i>Dst</i> occurred	Solar Radio Flux	Sunspot number	Year	No. of <i>Dst</i> occurred	Solar Radio Flux	Sunspot number
1996	6	72.0	12	2011	12	73.1	81
1997	19	80.9	29	2012	23	69.0	85
1998	22	118.1	88	2013	22	122.7	94
1999	17	153.9	136	2014	13	145.9	113
2000	31	180.0	174	2015	27	117.7	70
2001	19	181.1	170	2016	20	88.7	40
2002	18	179.4	164	2017	16	77.3	22
2003	25	128.4	99	2018	7	69.9	7
2004	10	106.8	65	2019	6	69.7	4
2005	17	91.7	46	2020	3	73.7	9
2006	10	80.0	25	2021	9	81.6	30
2007	4	73.1	13	2022	18	125.0	83
2008	4	69.0	4	2023	23	159.7	125
2009	1	70.5	5	2024	19	188.2	155
2010	8	80.0	25				

Figure 1 explains the following points:

- **SC23** (1996–2008): Shows a strong peak in solar activity around 2000, as indicated by high values of both solar radio flux and sunspot numbers. The number of *Dst* occurrences also peaks during this period, indicating a higher frequency of geomagnetic storms.
- **SC24** (2008–2019): The peak in solar activity is less pronounced compared to SC23, with both solar flux and sunspot numbers showing lower maximum values around 2014. Correspondingly, the number of *Dst* occurrences also appears lower.
- **SC25** (2019–2024): The current cycle shows an upward trend in solar activity, with solar flux and sunspot numbers increasing, and *Dst* occurrences starting to rise.

Between each solar cycle, there are periods of low solar activity (solar mini-




Figure 1. Correlation between the three parameters of three solar cycles i.e., solar radio flux, sunspot number and the occurrences of the geomagnetic storm.

mums). The most significant drop-in activity can be seen around 2008, which marks the transition from SC23 to SC24.

From Figure 1, we can conclude that the rising phase of solar cycle 25 is closely aligned with the behavior observed in solar cycle 23, suggesting similar patterns of solar activity. In contrast, solar cycle 24 stands out as being distinct from both cycles, exhibiting lower solar activity overall. Based on this comparison, it can be inferred that solar cycle 25 may follow a trajectory similar to that of solar cycle 23.

The graph, i.e. Figure 2 illustrates the cyclical nature of solar activity and its direct impact on geomagnetic storm occurrences. As solar radio flux increases, geomagnetic storm occurrences tend to rise, indicating a clear link between solar activity and geomagnetic disturbances on Earth. SC25 shows early signs of mimicking the behavior of SC23, suggesting that the upcoming peak in solar activity may bring a higher frequency of geomagnetic storms, similar to the intense activity seen during SC23.

The graph, i.e. Figure 3 demonstrates the cyclical nature of solar activity, with peaks in both sunspot numbers and solar radio flux during the maximum phases of each solar cycle. Solar cycle 24 stands out as being weaker than the adjacent




Figure 2. Comparison between the yearly averaged solar radio flux with the number of occurrences of geomagnetic storms from 1996 to 2024, covering solar cycles 23, 24 and the rising phase of solar cycle 25.




Figure 3. Comparison between the yearly averaged sunspot number and yearly averaged solar radio flux from 1996 to 2024, encompassing solar cycles 23, 24 and rising phase of solar cycle 25.




Figure 4. Comparison between the yearly averaged sunspot number and the number of occurrences of geomagnetic storms from 1996 to 2024, covering solar cycles 23, 24 and the rising phase of solar cycle 25.

cycles, while solar cycle 25 is on track to reach or exceed the levels of solar activity observed during solar cycle 23. This strong rise in solar activity during SC25 is crucial for understanding its potential impact on space weather and geomagnetic disturbances.

The graph, i.e., Figure 4 highlights the relationship between solar activity (as indicated by sunspot numbers) and geomagnetic storm occurrences. Solar cycles 23 and 25 show a strong correlation between increased sunspot numbers and geomagnetic storms, while solar cycle 24 stands out as a period of reduced activity. The rising phase of solar cycle 25 suggests that it could experience solar activity levels similar to those seen in solar cycle 23, which may lead to an increase in geomagnetic storm occurrences in the coming years.

The three charts, i.e. Figures 5–7 together illustrate the interconnectedness between solar activity indicators, such as solar flux and sunspot numbers, and the occurrences of geomagnetic storms. The peaks in solar flux and sunspot numbers align with periods of heightened geomagnetic storm occurrences. Solar Cycle 25 is exhibiting patterns similar to those of Solar Cycle 23, suggesting that the coming years may experience heightened solar activity and corresponding geomagnetic disturbances, compared to the weaker Solar Cycle 24. This




Figure 5. No. of occurrences of geomagnetic storm from the year 1996 to 2024.

reinforces the importance of understanding solar cycles for better forecasting of geomagnetic storms, which have significant impacts on Earth’s technological infrastructure.




Figure 6. Yearly averaged solar radio flux from the year 1996 to 2024.




Figure 7. yearly averaged sunspot number from the year 1996 to 2024.

### 3 Result

After analyzing and observing solar activities and geomagnetic field disturbances from 1996 to 2024, we identified a total of 431 geomagnetic storms. The data suggests a clear dependence between solar activity – such as sunspot number and solar radio flux—and the occurrence of geomagnetic storms. Specifically, the study revealed a positive correlation of 0.70 between the yearly average sunspot number and geomagnetic storm occurrences, a positive correlation of 0.61 between yearly average solar radio flux and storm occurrences, and a strong correlation of 0.93 between sunspot number and solar radio flux.

### 4 Conclusion

This paper has examined the interrelationships between key solar activity indicators – sunspot number, solar radio flux, and the Disturbance Storm Time (*Dst*) index – highlighting their vital roles in understanding solar variability and its impacts on Earth’s magnetosphere and ionosphere. These parameters are not only central to characterizing solar activity but also crucial for enhancing space weather forecasting capabilities. By reviewing the historical trends and significance of these indicators, we have emphasized their influence on major space weather phenomena such as geomagnetic storms and solar radiation storms.

Our analysis of various methodologies, including traditional statistical models and emerging machine learning techniques, provides a comprehensive approach to predicting solar activity. By elucidating the correlations and interactions between sunspot numbers, solar radio flux, and the *Dst* index, this study offers

new insights into solar-terrestrial relationships. These findings have the potential to improve the accuracy of space weather predictions, thereby aiding in the development of more effective mitigation strategies to minimize the impact of solar events on critical technological infrastructure.

In conclusion, this research contributes to the broader understanding of space weather science by advancing our knowledge of how solar activity indicators interact with geomagnetic events. As solar variability continues to pose challenges for Earth's technological systems, this work serves as a step forward in the ongoing effort to enhance space weather forecasting and safeguard modern technology from the adverse effects of solar activity.

### Acknowledgement

The authors would like to acknowledge the use of the OMNI data obtained from the GSFC/SPDF OMNI Web interface at <https://omniweb.gsfc.nasa.gov>. The data utilized in this research, including Sunspot Number, Solar Radio Flux, and Disturbance Storm Time (*Dst*) Index, have been invaluable to the study and analysis presented in this paper. We extend our gratitude to the Space Physics Data Facility (SPDF) for providing this critical resource.

### References

- [1] S. Yamakawa, M. Inoue, R. Suppiah (2016) Relationships between solar activity and variations in SST and atmospheric circulation in the stratosphere and troposphere. *Quat. Int.* **397** 289-299.
- [2] B. Van. Geel, S. Engels (2012) The effects of changing solar activity on climate: contributions from palaeoclimatological studies. *J. Space Weather Space Clim.* **2** (9).
- [3] R. Sharma, O.P. Tripathi, A. Dwivedi, S. Kumar (2023) Association of sunspot number, solar radio flux and geomagnetic storms during the period of 2009-2019. *Int. J. Recent Sci. Res.* **14** (6).
- [4] B. Adhikari, S. Dahal, R.K. Mishra, N. Sapkota, et al. (2019) Analysis of solar, interplanetary, and geomagnetic parameters during solar cycles 22, 23 and 24. *Russ. J. Earth Sci.* **19** ES1003.
- [5] D.H. Hathway (2010) The Solar Cycle. *Living Rev. Sol. Phys.* **7**(1) (2010).
- [6] R.G. Cionco, S.M. Kudryavtsev, W.W-H. Soon (2023) Tidal Forcing on the Sun and the 11-Year Solar-Activity Cycle. *Solar Phys.* **298**(5) 70.
- [7] R. Schlichenmaier, R. Rezaei, G.N. Bello, T. A. Waldmann (2010) The formation of a sunspot penumbra. *Astron. Astrophys.* **512** L1.
- [8] I. Kutiev, I. Tsagouri, L. Perrone, D. Pancheva, P. Mukhtarov, A. Mikhailov, J. Lastovicka, N. Jakowski, D. Buresova, E. Blanch, B. Andonov, D. Altadill, S. Magdaleno, M. Parisi, J.M. Tort (2013) Solar activity impact on the Earth's upper atmosphere. *J. Space Weather Space Clim.* bf3 (21).
- [9] N.V. Nitta, T. Mulligan, E.K.J. Kilpua, B.J. Lynch, M. Mierla, J.O. Kane, P. Pagano, E. Palmerio, J. Pomoell, G.I. Richardson, L. Rodriguez, A.P. Rouillard, S. Sinha,

- N. Srivastava, D.-C. Talpeanu, S.L. Yardley, A.N. Zhukov (2021) Understanding the Origins of Problem Geomagnetic Storms Associated with “Stealth” Coronal Mass Ejections. *Space Sci. Rev.* **217**(8) 82.
- [10] G.V. Kuklin (1976) *Contrib. Astron. Obs. Skalnaté Pleso* **6** 195.
- [11] E. Saiz, Y. Cerrato, C. Cid, V. Dobrica, P. Hejda, P. Nenovski, P. Stauning, J. Bochnicek, D. Danov, C. Demetrescu, W.D. Gonzalez, G. Maris, D. Teodosiev, F. Valach (2013) Geomagnetic response to solar and interplanetary disturbances. *J. Space Weather Space Clim.* **3** (20).
- [12] G.S. Lakhina, B.T. Tsurutani (2016) Geomagnetic storms: historical perspective to modern view. *Geosci. Lett.* **3** 5.
- [13] J. Feynman, N.U. Crooker (1978) The solar wind at the turn of the century. *Nature* **275** 626–627.
- [14] L. Svalgaard et al. (1978) The strength of the Sun’s polar fields. *Solar Phys.* **58** 225–240.
- [15] E.W. Cliver, V. Boriakoff, K.H. Bounar (1996) The 22-year cycle of geomagnetic and solar wind activity. *J. Geophys. Res. Space Physics* **101** 27091–27109.
- [16] ??? M. Lockwood et al. (1999) *Nature* **399** 437–439.
- [17] ??? R. Stamper, M. Lockwood, M.N. Wild, T.D.G. Clark (1999) Solar Causes of the Long-Term Increase in Geomagnetic Activity. *J. Geophys. Res. Space Physics* **104** 325–342.
- [18] O.M. Raspopov, V.A. Dergachev, J. Esper, O. Kozyreva (2008) The influence of the de Vries (~200-year) solar cycle on climate variations: Results from the Central Asian Mountains and their global link. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **259**(1) 6–16.
- [19] K.F. Tapping (2013) The 10.7 cm solar radio flux (F10.7). *Space Weather* **11**(7) 394–406.
- [20] G.S. Lakhina (1994) Solar wind-magnetosphere-ionosphere coupling and chaotic dynamic. *Surv. Geophys.* **15** 703–754.
- [21] J.E. Borovsky, Y.Y. Shprits (2017) Is the Dst Index Sufficient to Define All Geospace Storms? Is Dst Sufficient to Define Storms? *J. Geophys. Res. Space Physics* **122** (3).
- [22] X. Meng, B.T. Tsurutani, A.J. Manucci (2019) The Solar and Interplanetary Causes of Superstorms (Minimum Dst  $\leq -250$  nT) During the Space Age. *J. Geophys. Res. Space Physics* **124** 3926–3948.
- [23] H. Nemethova, J. Zyka (2024) Overview of the Space Weather Impact on the Navigation and Communication Systems. *Acta Avion.* **26**(1) 27–39.
- [24] N. Buzulukova, B. Tsurutani (2022) Space Weather: From solar origins to risks and hazards evolving in time. *Front. Astron. Space Sci.* **9**.